


NEW ROOTS

NATURAL HEALTH PRODUCTS
WWW.NEWROOTSHERBAL.EU

HERBAL

» WOMEN'S HEALTH



- Exclusive information for health-care professionals -



FEMINA RELIEF

SYNERGISTIC COMBINATION OF 8 PLANTS
HIGH POTENCY STANDARDISED EXTRACTS

PREMENSTRUAL SYNDROME

EMOTIONAL SYMPTOMS

- » Irritability
- » Mood swings
- » Lack of concentration
- » Depression
- » Lack of energy
- » Changes in appetite (cravings)

PHYSICAL SYMPTOMS

- » Breast pain (mastalgia)
- » Generalised aches and pains (arthralgias and myalgias)
- » Headaches
- » Abdominal distention
- » Fluid retention



Product Datasheet

90 capsules Code FE0253

COMPREHENSIVE FORMULA FOR ACTION ON ALL SYMPTOMS

Nutritional information:	1 capsule
Dandelion root ⁽¹⁾ (3% flavonoids) (10:1)	150 mg
Chasteberry ⁽¹⁾ (0.5% agnuside) (25:1)	125 mg
Dong quai ⁽¹⁾ (1% Z-ligustilide) (20:1)	125 mg
Cramp bark ⁽¹⁾ (4:1)	50 mg
Corn silk ⁽¹⁾ (4:1)	50 mg
Red clover ⁽¹⁾ (8% isoflavones)	35 mg
Black cohosh ⁽¹⁾ (2,5% triterpene glycosides) (12:1)	20 mg
Siberian ginseng ⁽¹⁾ (0,8% eleutherosides) (50:1)	5 mg

⁽¹⁾ standardised extracts

Recommended daily dose: 1 capsule daily with food.

References:

1. Kupfersztain, C., et al. "The immediate effect of natural plant extract, *Angelica sinensis* and *Matricaria chamomilla* (Climex) for the treatment of hot flushes during menopause. A preliminary report." *Clinical and experimental obstetrics & gynecology* 30.4 (2002): 203-206.
2. Mirghafourvand, Mojgan, et al. "Effects of *Vitex agnus* and Flaxseed on cyclic mastalgia: A randomized controlled trial." *Complementary therapies in medicine* 24 (2016): 90-95.
3. Ambrosini, Anna, et al. "Use of *Vitex agnus-castus* in migrainous women with premenstrual syndrome: an open-label clinical observation." *Acta Neurologica Belgica* 113.1 (2013): 25-29.
4. He, Zhong, et al. "Treatment for premenstrual syndrome with *Vitex agnus castus*: A prospective, randomized, multi-center placebo controlled study in China." *Maturitas* 63.1 (2009): 99-103.
5. Tanmahasamut, P., et al. "*Cimicifuga racemosa* extract for relieving menopausal symptoms: a randomized controlled trial." *Climacteric* 18.1 (2015): 79-85.
6. Samman, Samir, et al. "Red clover (*Trifolium pratense*) isoflavones and serum homocysteine in premenopausal women: a pilot study." *Journal of Women's Health* 18.11 (2009): 1813-1816.
7. Blakesmith, Sarah J., et al. "Effects of supplementation with purified red clover (*Trifolium pratense*) isoflavones on plasma lipids and insulin resistance in healthy premenopausal women." *British Journal of Nutrition* 89.04 (2003): 467-474.
8. Clare, Bevin A., Richard S. Conroy, and Kevin Spelman. "The diuretic effect in human subjects of an extract of *Taraxacum officinale* folium over a single day." *The Journal of Alternative and Complementary Medicine* 15.8 (2009): 929-934.
9. Doan, Du Dat, et al. "Studies on the individual and combined diuretic effects of four Vietnamese traditional herbal remedies (*Zea mays*, *Imperata cylindrica*, *Plantago major* and *Orthosiphon stamineus*)." *Journal of ethnopharmacology* 36.3 (1992): 225-231.
10. Pellow J. The Efficacy of *Viburnum opulus* 3X in the Treatment of Primary Dysmenorrhea. 2016. (<https://clinicaltrials.gov/ct2/show/NCT02467543>)
11. Bohn, B., C. T. Nebe, and C. Birr. "Flow-cytometric studies with *Eleutherococcus senticosus* extract as an immunomodulatory agent." *Arzneimittel-Forschung* 37.10 (1987): 1193-1196.
12. Hartz, A. J., et al. "Randomized controlled trial of Siberian ginseng for chronic fatigue." *Psychological medicine* 34.01 (2004): 51-61.

DONG QUAI (*Angelica sinensis*) ⁽¹⁾

- » Female tonic
- » Dysmenorrhoea, general discomfort and bloating
- » Mood swings

*CHASTEBERRY (*Vitex agnus-castus*) ⁽²⁻⁴⁾

- » Female hormonal balance
- » Breast pain (mastalgia), swelling, retention, anxiety, irritability, mood swings

*BLACK COHOSH (*Cimicifuga racemosa*) ⁽⁵⁾

- » Uterine cramps
- » Depression, anxiety, mood swings

CORN SILK (*Zea mays*) ⁽⁹⁾

- » Fluid retention
- » Urinary tract infections

DANDELION (*Taraxacum officinale*) ⁽⁸⁾

- » Fluid retention
- » Hepatic detoxification oestrogen elimination

CRAMP BARK (*Viburnum opulus*) ⁽¹⁰⁾

- » Diuretic
- » Spasmodic effect for menstrual cramps

SIBERIAN GINSENG

(*Eleutherococcus senticosus*) ⁽¹¹⁻¹²⁾

- » General tonic
- » Adrenal support for dysfunction (stress, depression, fatigue and irritability)

RED CLOVER (*Trifolium pratense*) ⁽⁶⁻⁷⁾

- » Oestrogen modulator

*E Commission Approved: Therapeutic Guide to Medicinal Plants of the German Ministry of Health



FEMINA FLORA ORAL ORAL ADMINISTRATION OF LACTOBACILLUS REACHES THE VAGINAL MUCOSA WITHIN ONE WEEK (12-18)

NATURAL ALTERNATIVE FOR VAGINAL CANDIDIASIS PROBIOTIC FOR A BALANCED VAGINAL FLORA

GENITOURINARY INFECTIONS

- » They are the most common reason for gynaecological consultations.
- » They can be bacterial in nature (bacterial vaginosis) or fungal (vulvovaginal candidiasis).
- » The types of bacteria in the vaginal canal vary depending on hormone levels, diet, sexual contact and douching.
- » *Lactobacillus* are the dominant bacteria of the vaginal flora, changes in this vaginal flora can lead to urinary and vaginal infections ⁽¹⁾.
- » Some probiotics can help modulate the immune system and help displace the pathogens that cause these infections ⁽²⁾.
- » Lactobacillus produce H₂O₂ helping to maintain an optimal vaginal pH close to 4.0 and decreasing the risk of developing vaginosis ⁽³⁾.



Product Datasheet

GPS™ capsule

16 strains
55,000 Mill. CFU
With FOS and AOS

Dose: 1 to 2 capsules a day

Format: 30 capsules

PREVENTION OF VAGINAL INFECTIONS IDEAL FOR PROLONGED USE

Code FE2286


It can help prevent:

- » **Recurrent urinary tract infections**⁽¹⁾.
- » **Bacterial vaginosis** ⁽²⁻⁴⁾ and recurrent vulvovaginal candidiasis ⁽⁵⁻⁹⁾.
- » **Group B Streptococcus (GBS)** growth during pregnancy ^(10,11).

Nutritional information: 1 capsule

Human strains:	
<i>Lactobacillus rhamnosus</i> UB5115	19,000 billion CFU
<i>Lactobacillus acidophilus</i> UB5997	5,250 billion CFU
<i>Lactobacillus casei</i> UB1499	12,000 billion CFU
<i>Bifidobacterium bifidum</i> UB4280	2,000 billion CFU
<i>Bifidobacterium breve</i> UB8674	2,000 billion CFU
<i>Bifidobacterium longum</i> UB7691	2,000 billion CFU
<i>Lactobacillus crispatus</i> UB4719	0,750 billion CFU
<i>Lactobacillus gasserii</i> UB8141	0,750 billion CFU
<i>Lactobacillus rhamnosus</i> GG	0,300 billion CFU
<i>Lactobacillus acidophilus</i> LA-14	0,300 billion CFU
Dairy strains:	
<i>Lactobacillus casei</i> LC-11	0,300 billion CFU
<i>Lactobacillus helveticus</i> UB7229	0,500 billion CFU
<i>Lactobacillus paracasei</i> UB1978	0,600 billion CFU
<i>Lactobacillus johnsonii</i> UB3394	0,750 billion CFU
<i>Lactobacillus reuteri</i> UB2419	0,500 billion CFU
Plant strain:	
<i>Lactobacillus plantarum</i> UB2783	8,000 billion CFU
Inulin	10 mg
Arabinogalactan (AOS)	10 mg

PREVENTION OF VAGINAL CANDIDIASIS AND BACTERIAL VAGINITIS




References:

1. Stapleton, Ann E., et al. "Randomized, placebo-controlled phase 2 trial of a *Lactobacillus crispatus* probiotic given intravaginally for prevention of recurrent urinary tract infection." *Clinical infectious diseases* 52.10 (2011): 1212-1217.
2. Delia, A., et al. "Effectiveness of oral administration of *Lactobacillus paracasei* subsp. *paracasei* F19 in association with vaginal suppositories of *Lactobacillus acidophilus* in the treatment of vaginosis and in the prevention of recurrent vaginitis." *Minerva ginecologica* 58.3 (2006): 227-231.
3. Marcone, Valentina, et al. "Long term vaginal administration of *Lactobacillus rhamnosus* as a complementary approach to management of bacterial vaginosis." *International Journal of Gynecology & Obstetrics* 110.3 (2010): 223-226.
4. Ya, Wang, Cheryl Reifer, and Larry E. Miller. "Efficacy of vaginal probiotic capsules for recurrent bacterial vaginosis: a double-blind, randomized, placebo-controlled study." *American journal of obstetrics and gynecology* 203.2 (2010): 120-e1.
5. Coudeyras, Sophie, et al. "Adhesion of human probiotic *Lactobacillus rhamnosus* to cervical and vaginal cells and interaction with vaginosis and vaginitis-associated pathogens." *Infectious Diseases in Obstetrics and Gynecology* 2008 (2008): 1.
6. Falagas, Matthew E., Gregoria I. Betsi, and Stavros Athanasiou. "Probiotics for prevention of recurrent vulvovaginal candidiasis: a review." *Journal of Antimicrobial Chemotherapy* 58.2 (2006): 266-272.
7. Li, Duo, et al. "Efficacy and safety of probiotics in the treatment of *Candida* associated stomatitis." *Mycoses* 57.3 (2014): 141-146.
8. Pirotta, Marie, et al. "Effect of *Lactobacillus* in preventing post-antibiotic vulvovaginal candidiasis: a randomised controlled trial." *BMJ* 329.7465 (2004): 548.
9. Strus, M., et al. "Inhibitory activity of vaginal *Lactobacillus* bacteria on yeasts causing vulvovaginal candidiasis." *Medycyna Do wadczalna* 57.1 (2005): 16.
10. Zarate, G., and M. E. Nader Macias. "Influence of probiotic vaginal *Lactobacilli* on in vitro adhesion of urogenital pathogens to vaginal epithelial cells." *Letters in applied microbiology* 43.2 (2006): 174-180.
11. Açıköz, Z. C., et al. "Inhibitor effect of vaginal *Lactobacilli* on group B streptococci." *Mikrobiyoloji bulteni* 39.1 (2005): 17-23.
12. Reid, Gregor, et al. "Oral use of *Lactobacillus rhamnosus* GR-1 and *L. fermentum* RC-14 significantly alters vaginal flora: randomized, placebo-controlled trial in 64 healthy women." *FEMS Immunology & Medical Microbiology* 35.2 (2003): 131-134.
13. Petricevic, Ljubomir, et al. "Randomized, double-blind, placebo-controlled study of oral *Lactobacilli* to improve the vaginal flora of postmenopausal women." *European Journal of Obstetrics & Gynecology and Reproductive Biology* 141.1 (2008): 54-57.
14. Grewal, Nipunjot, Amita Mahajan, and Jagminder Kaur Bajaj. "The Effect of Supplementation of Standard Antibiotic Therapy with Oral Probiotics for Bacterial Vaginosis." *International Journal of Medical and Dental Sciences* 71 (2018): 1628-1631.
15. Reid, Gregor, et al. "Oral probiotics can resolve urogenital infections." *FEMS Immunology & Medical Microbiology* 30.1 (2001): 49-52.
16. Strus, Magdalena, et al. "Studies on the effects of probiotic *Lactobacillus* mixture given orally on vaginal and rectal colonization and on parameters of vaginal health in women with intermediate vaginal flora." *European Journal of Obstetrics & Gynecology and Reproductive Biology* 163.2 (2012): 210-215.
17. Hilton, Eileen, et al. "Ingestion of yogurt containing *Lactobacillus acidophilus* as prophylaxis for candidal vaginitis." *Annals of Internal Medicine* 116.5 (1992): 353-357.
18. Reid, Gregor, et al. "Probiotic *Lactobacillus* dose required to restore and maintain a normal vaginal flora." *FEMS Immunology & Medical Microbiology* 32.1 (2001): 37-41.

D-MANOCIST

TRIPLE ACTION FOR THE WELL-BEING OF THE URINARY TRACT⁽¹⁾



Nutritional information:	2 scoops	Recommended daily dose: Mix 2 scoops (approx. 5,4 g) in water or juice twice daily for two days, then 2 scoops (approx. 5,4 g) in water or juice once daily.
D-mannose	4.800 mg	
Cranberry 107:1	600 mg	
<i>Lactobacillus rhamnosus</i> R0011	500 mill. CFU	

- » **D-MANNOSE** ⁽²⁻⁵⁾. Natural sugar that reaches the source of infection directly, without metabolism, and prevents the adhesion of *E. coli type 1 fimbriae* on the bladder wall so that the bacteria will be washed away with the urine flow.
- » **CRANBERRY** (*V. macrocarpon*)⁽³⁻⁸⁾. *A-type* proanthocyanidins (A PACs), active constituents of cranberry, prevent the adhesion of *E. coli* P-type fimbriae to uroepithelial cells, thus preventing their colonisation.
- » **PROBIOTICS** (*Lactobacillus rhamnosus*)⁽⁹⁻¹²⁾. Lactobacilli are dominant in the vaginal ecosystem and the rectum indirectly acts as a reservoir for vaginal lactobacilli. Its protective role is focused:
 - » They interfere with the adherence of uropathogens to the vaginal epithelium.
 - » They produce antimicrobial compounds (organic acids, H₂O₂ and bacteriocins) and create an environment of vaginal acidity (pH 4-4.5) that prevents pathogens from establishing themselves.
 - » Coaggregation of pathogens to maximise antimicrobial effect.
 - » Amplification of the immune response by *Lactobacillus* strains.

References:

1. Vicariotto, Franco. "Effectiveness of an association of a cranberry dry extract, D-mannose, and the two microorganisms *Lactobacillus plantarum* LP01 and *Lactobacillus paracasei* LPC09 in women affected by cystitis: a pilot study." *Journal of clinical gastroenterology* 48 (2014): S96-S101.
2. Altarac, Silvio, and Dino Papeš. "Use of d mannose in prophylaxis of recurrent urinary tract infections (UTIs) in women." *BJU international* 113.1 (2014): 9-10.
3. Kranj ec, Bojana, Dino Papeš, and Silvio Altarac. "D-mannose powder for prophylaxis of recurrent urinary tract infections in women: a randomized clinical trial." *World journal of urology* 32.1 (2014): 79-84.
4. Wright, J. V. "D-Mannose for bladder and kidney infections." *Townsend Letter for Doctors and Patients* 192 (1999): 96-8.
5. Ofek, I, A. Mosek, and N. Sharon. "Mannose-specific adherence of *Escherichia coli* freshly excreted in the urine of patients with urinary tract infections, and of isolates subcultured from the infected urine." *Infection and immunity* 34.3 (1981): 708-711.
6. Foxman, Betsy, et al. "Cranberry juice capsules and urinary tract infection after surgery: results of a randomized trial." *American journal of obstetrics and gynecology* 213.2 (2015): 194-e1.
7. Kaspar, Kerrie L., Amy B. Howell, and Christina Khoo. "A randomized, double-blind, placebo-controlled trial to assess the bacterial anti-adhesion effects of cranberry extract beverages." *Food & function* 6.4 (2015): 1212-1217.
8. Ledda, A., et al. "Cranberry supplementation in the prevention of non-severe lower urinary tract infections: a pilot study." *Eur Rev Med Pharmacol Sci* 19.1 (2015): 77-80.
9. Shim, Yoon Hee, Seung Joo Lee, and Jung Won Lee. "Antimicrobial activity of lactobacillus strains against uropathogens." *Pediatrics International* 58.10 (2016): 1009-1013.
10. Martín, Rebeca, et al. "La microbiota vaginal: composición, papel protector, patología asociada y perspectivas terapéuticas." *Enfermedades Infecciosas y Microbiología Clínica* 26.3 (2008): 160-167.
11. Beerepoot, M. A. J., Gerben Ter Riet, and Sita Nys. "Lactobacilli versus antibiotics to prevent urinary tract infections. A randomized double-blind non-inferiority trial in postmenopausal women." *Archives of internal medicine* 172.9 (2012): 704-712.
12. Grin, Peter M., et al. "Lactobacillus for preventing recurrent urinary tract infections in women: meta-analysis." *Can J Urol* 20.1 (2013): 6607-6614.

OBSERVATIONAL TRIAL

"Natural treatment of recurrent cystitis with D-Manocist"


Study of the efficacy and effectiveness of "D-Manocist" in 20 women with recurrent cystitis, "Plan de Farmacias Diferentes" (Nafarco).

Treatment: 3 teaspoons twice a day for 2 days. 3 teaspoons, once a day until the first bottle is finished. 7 days off. 1 teaspoon once a day for 10 days.

CONCLUSIONS

1. Reduction of symptomatology in 90% of the population.
2. Improvement from the 1st week of treatment.
3. 84% of the population had no recurrence.
4. 100% would take the product again and/or recommend it.


Symptomatology evolution of the population





INOSITOL (*myo*-inositol)

POLYCYSTIC OVARY SYNDROME / FEMALE INFERTILITY



Product Datasheet

125 gr. Code: FE2208

- » Considered a member of the B vitamin group (Bh or B8).
- » It has benefits for blood sugar regulation, as well as for mood.
- » It plays a very important role in fertility.
- » Integral component of healthy cell membranes, high resistance to oxidative stress.

Nutritional information: 1 scoop (4 g)

Inositol (*myo*-inositol) 4 g

Recommended daily dose: 1 scoop (4 g) added to 250 ml of water or juice, once daily. Consult a health-care practitioner for use beyond 6 weeks.

Manufactured under the strict quality standards of the United States Pharmacopeia
(United States Pharmacopeia, USP)

POLYCYSTIC OVARY SYNDROME (PCOS)⁽¹⁾

PCOS is a common cause of infertility, with symptoms including irregular menstruation, failure to release an egg during menstruation (anovulation), ovarian cysts and unwanted hair growth with a systemic male pattern.

Inositol reduces oxidative stress, restores ovulation and oocyte quality, normalises the menstrual cycle and reduces serum testosterone in women with PCOS.



FEMALE FERTILITY / IN VITRO FERTILISATION⁽²⁾

It improves embryo and oocyte quality. It can be an alternative to prescription medications, alone or as part of assisted reproductive therapy.



Other benefits:

- » **Gestational diabetes**⁽³⁾: reduces risk and improves insulin resistance.
- » **Metabolic syndrome**⁽⁴⁾: improves blood pressure, insulin resistance, and cholesterol and triglyceride levels.
- » **Premenstrual dysphoric disorder (PMDD)**⁽⁵⁾: improves symptoms.
- » **Depression**⁽⁶⁾: improves symptoms of depression even in people with bipolar disorder.
- » **Panic disorder**⁽⁷⁾: reduces the frequency and severity of attacks, as well as the severity of agoraphobia.
- » **Obsessive compulsive disorder**⁽⁸⁻⁹⁾: is effective in its treatment as well as in trichotillomania (obsessive disorder of pulling out one's own hair).
- » **Bulimia**⁽¹⁰⁾: improves the symptoms of eating disorders.
- » **Chronic thyroiditis (Hashimoto's disease)**⁽¹¹⁾: improves thyroid function.
- » **Cancer**⁽¹²⁾: as an adjuvant, it reduces the adverse effects of chemotherapy.
- » **Neural tube defects**⁽¹³⁾: may help prevent folic acid-resistant neural tube defects.
- » **Male fertility**⁽¹⁴⁾: improves sperm quality (concentration, motility, total sperm count, etc.) and regulates hormone levels (LH, FSH and Inhibin B).

References:

1. Fruzzetti, Franca, et al. "Comparison of two insulin sensitizers, metformin and *myo*-inositol, in women with polycystic ovary syndrome (PCOS)." *Gynecological Endocrinology* 33.1 (2017): 39-42.
2. Caprio, Francesca, et al. "Myo-inositol therapy for poor-responders during IVF: a prospective controlled observational trial." *Journal of ovarian research* 8.1 (2015): 37.
3. Santamaria, Angelo, et al. "Myo-inositol may prevent gestational diabetes onset in overweight women: a randomized, controlled trial." *The Journal of Maternal-Fetal & Neonatal Medicine* 29.19 (2016): 3234-3237.
4. Giordano, Domenico, et al. "Effects of *myo*-inositol supplementation in postmenopausal women with metabolic syndrome: a perspective, randomized, placebo-controlled study." *Menopause* 18.1 (2011): 102-104.
5. Carlomagno, Gianfranco, et al. "Myo inositol in the treatment of premenstrual dysphoric disorder." *Human Psychopharmacology: Clinical and Experimental* 26.7 (2011): 526-530.
6. Levine, Joseph, et al. "Double-blind, controlled trial of inositol treatment of depression." *The American journal of psychiatry* 152.5 (1995): 792.
7. Palatnik, Alex, et al. "Double-blind, controlled, crossover trial of inositol versus fluvoxamine for the treatment of panic disorder." *Journal of clinical psychopharmacology* 21.3 (2001): 335-339.
8. Fux, Mendel, et al. "Inositol treatment of obsessive-compulsive disorder." *The American journal of psychiatry* 153.9 (1996): 1219.
9. Leppink, Eric W., Sarah A. Redden, and Jon E. Grant. "A double-blind, placebo-controlled study of inositol in trichotillomania." *International clinical psychopharmacology* 32.2 (2017): 107-114.
10. Gelber, Diana, Joseph Levine, and R. H. Belmaker. "Effect of inositol on bulimia nervosa and binge eating." *International Journal of Eating Disorders* 29.3 (2001): 345-348.
11. Nordio, M., and S. Basciani. "Myo-inositol plus selenium supplementation restores euthyroid state in Hashimoto's patients with subclinical hypothyroidism." *Eur Rev Med Pharmacol Sci* 21.2 Suppl (2017): 51-59.
12. Bai, Ivan, et al. "Efficacy of IP 6+ inositol in the treatment of breast cancer patients receiving chemotherapy: prospective, randomized, pilot clinical study." *Journal of Experimental & Clinical Cancer Research* 29.1 (2010): 12.
13. Greene, Nicholas DE, et al. "Inositol for the prevention of neural tube defects: a pilot randomised controlled trial." *British Journal of Nutrition* 115.6 (2016): 974-983.
14. Calogero, A. E., et al. "Myoinositol improves sperm parameters and serum reproductive hormones in patients with idiopathic infertility: a prospective double blind randomized placebo controlled study." *Andrology* 3.3 (2015): 491-495.

NATAL NUTRI BALANCE

THE MOST COMPLETE FORMULA TO ENSURE NUTRITIONAL DEMANDS ARE MET DURING PREGNANCY AND LACTATION



Product Datasheet

90 capsules Code FE1434

23 VITAMINS AND MINERALS

In their most bioavailable forms

Nutritional information:	1 capsule
Folate (calcium-L-methylfolate)	333,3 µg (167%*)
Thiamin (vit. B1) (from 25 mg thiamin hydrochloride)	22,3 mg (2 027%*)
Riboflavin (vit. B2) (from 11,67 mg riboflavin-5-phosphate sodium)	8,8 mg (629%*)
Niacin (vit. B3) (from nicotinamide)	16,7 mg NE (104%*)
Pantothenic acid (vit. B5) (from 33,33 mg d-pantothenate calcium)	30,5 mg (508%*)
Vitamin B6 (from 11,67 mg pyridoxal-5-phosphate)	8 mg (571%*)
Vitamin B12 (methylcobalamin)	333,3 µg (13 332%*)
Biotin (D-biotin)	100 µg (200%*)
Vitamin C (from 58,33 mg calcium-L-ascorbate)	47,9 mg (60%*)
Vitamine D3 (from cholecalciferol) (133,3 IU/capsule)	3,3 µg (66%*)
Vitamine E (from d-alpha-tocopherol) (16,7 IU/capsule)	12,3 mg α-TE (103%*)
Vitamine K1 (phyloquinone)	33,3 µg (44%*)
Manganese (from manganese citrate)	1,7 mg (85%*)
Iron (from iron bisglycinate)	11,7 mg (84%*)
Zinc (from zinc citrate)	8,3 mg (83%*)
Chromium (from chromium picolinate)	33,3 µg (83%*)
Copper (from copper citrate)	0,5 mg (50%*)
Iodine (from potassium iodide)	50 µg (33%*)
Molybdenum (from sodium molybdate)	16,7 µg (33%*)
Selenium (from L-selenomethionine))	16,7 µg (30%*)
Magnesium (from magnesium bisglycinate)	33,5 mg (9%*)
Calcium (from calcium citrate)	66,7 mg (8%*)
Potassium (from potassium citrate)	23,3 mg (1%*)

*NRV: Nutrient Reference Value in %

Recommended daily dose: 1 capsule daily with meals.

12

VITAMINS

Vitamin C, buffered for increased tolerability and bioavailability.

Vitamin D, essential for the baby's bone development.

Vitamin K1, involved in coagulation.

Vitamin E, deficiency of which is associated with rupture of the amniotic sac.

Complete B-complex, to avoid possible displacement.

11

MINERALS

Calcium, the foetus consumes 200-300 mg/day.

Iron, in chelated form for increased bioavailability.

Iodine, essential for a child's neurological development.

Magnesium, the need for which doubles in the third trimester. In its bisglycinate form for increased bioavailability.

Zinc, 82% of pregnant women are deficient. Risk of premature birth.

Chromium, deficiency is associated with gestational diabetes.

Selenium, copper, manganese, molybdenum and potassium, (other important minerals during pregnancy).



References:

- Liu, Jian-meng, et al. "Micronutrient supplementation and pregnancy outcomes: double-blind randomized controlled trial in China." *JAMA internal medicine* 173,4 (2013): 276-282.
- Kalra, Pramila, et al. "Effect of vitamin D supplementation during pregnancy on neonatal mineral homeostasis and anthropometry of the newborn and infant." *British Journal of Nutrition* 108.06 (2012): 1052-1058.
- Prado, Elizabeth L., et al. "Effects of maternal and child lipid-based nutrient supplements on infant development: a randomized trial in Malawi." *The American journal of clinical nutrition* 103.3 (2016): 784-793.
- Moeini, M. M., H. Karami, and E. Mikaeili. "Effect of selenium and vitamin E supplementation during the late pregnancy on reproductive indices and milk production in heifers." *Animal reproduction science* 114.1 (2009): 109-114.
- Zeng, Lingxia, et al. "Impact of micronutrient supplementation during pregnancy on birth weight, duration of gestation, and perinatal mortality in rural western China: double blind cluster randomised controlled trial." *Bmj* 337 (2008): a2001.
- Westphal, L. M., M. L. Polan, and A. Sontag Trant. "Double-blind, placebo-controlled study of Fertilityblend: a nutritional supplement for improving fertility in women." *Clinical and experimental obstetrics & gynecology* 33.4 (2005): 205-208.
- Correia-Santos, A. M., et al. "Dietary supplements for the lactating adolescent mother: influence on plasma micronutrients." *Nutr Hosp* 26.2 (2011): 392-398.
- Gupta, Piyush, et al. "Multimicronutrient supplementation for undernourished pregnant women and the birth size of their offspring: a double-blind, randomized, placebo-controlled trial." *Archives of pediatrics & adolescent medicine* 161.1 (2007): 58-64.
- Rush, D., et al. "The effects of dietary supplementation during pregnancy on placental morphology, pathology, and histomorphometry." *The American journal of clinical nutrition* 39.6 (1984): 863-871.

STRONG-BONES

Nutritional information:	2 capsules
Calcium (from hydroxyapatite)	300 mg (38%*)
Phosphorus (from hydroxyapatite)	150 mg (21%*)
Proteins (from hydroxyapatite)	300 mg
Magnesium (from Mg bisglycinate)	48,3 mg (**)
Zinc (from zinc mono-L-methionine sulfate)	3,1 mg (31%*)
Manganese (from Mn citrate)	0,9 mg (45%*)
Copper (from cupric citrate)	0,31 mg (31%*)
Boron (from boric acid)	1,1 mg
Field horsetail (<i>E. arvense</i>) (7% silica)	10 mg
Thiamin (vit. B1) (thiamin HCL)	1,4 mg (127%*)
Vit. K2 (menaquinone 4 and 7)	31 µg (41%*)
Vit. D3 (cholecalciferol, 167 IU/caps.)	8,3 µg (166%*)
Vit. C (L-ascorbic acid)	62 mg (78%*)
Vit. B12 (methylcobalamin)	50 µg (2 000%*)
Folate (calcium-L-methylfolate)	166,7 µg (83%*)
L-Lysine	100 mg
L-Proline	100 mg
Glucosamine sulfate (sodium-free)	84 mg
Turmeric (<i>C. longa</i>) (95% curcuminoids***)	6,7 mg
Grape seed (<i>V. vinifera</i>) (95% PACs)	20 mg
Green tea (<i>C. sinensis</i>) (5,25mg EGCG/caps.)	14 mg
Lutein (from <i>Tagetes erecta</i>)	0,7 mg
Lycopene (from <i>Lycopersicon esculentum</i>)	1,7 mg

* NRV: Nutrient Reference Value in %
 ** NRV: Nutrient Reference Value below 15%
 *** provides curcumin I, demetoxicurcumin and bisdemetoxicurcumin

Recommended daily dose: 2 capsules one to three times daily with food. If you are taking medications, take this product a few hours before or after taking them.

Microcrystalline Hydroxyapatite (MCH) (1-2) Source of calcium and osteospecific organic components (phosphorus, collagen and proteins) that contribute to stimulating osteogenesis. Optimum Calcium-Phosphorus ratio (2:1).


Oral administration of MCH is more effective than calcium carbonate in reducing trabecular bone loss in patients with osteoporosis. (3-4).



Product Datasheet

90 capsules Code FE1570
 180 capsules Code FE0297

MCH treatment VS calcium carbonate (CC) in osteoporosis.



Product Datasheet

180 capsules Code FE2084

Antioxidants: (6-10) exclusive antioxidant content with a very important role in bone formation (Lycopene, Lutein, Green Tea, Turmeric and Grape Seed). Protection against oxidative stress and regulation of inflammatory processes.

Magnesium Bisglycinate: (5) essential for Calcium-D3 metabolism. In bisglycinate form for 4 times higher bioavailability.

Co-Factors: (11-13) nutrient assimilation and bone formation. 6 Vitamins, 4 Minerals. Essential amino acids for the bone matrix (L-Proline and L-Lysine).

STRONG-BONES Vegetarian

Nutritional information:	3 capsules
Calcium (from calcium phosphate, tribasic)	333 mg (42%*)
Phosphorus (from calcium phosphate, tribasic)	157,4 mg (22%*)
Magnesium (from magnesium bisglycinate)	90 mg (24%*)
Zinc (from zinc mono-L-methionine sulphate)	3,1 mg (31%*)
Manganese (from manganese citrate)	0,93 mg (47%*)
Copper (from cupric citrate)	0,31 mg (31%*)
Boron (from boric acid)	1,1 mg
Field horsetail (<i>Equisetum arvense</i>) (7% silica)	10 mg
Thiamin (vit. B1) (from thiamin hydrochloride)	1,4 mg (127%*)
Vitamin K2 (from menaquinone-4 and menaquinone-7)	31 µg (41%*)
Vitamin D3 (cholecalciferol) (111 IU/caps.)	8,3 µg (166%*)
Vitamin C (L-ascorbic acid)	62 mg (78%*)
Vitamin B12 (methylcobalamin)	50 µg (2 000%*)
Folate (calcium-L-methylfolate)	166,7 µg (83%*)
L-Lysine	100 mg
L-Proline	100 mg
Glucosamine (from <i>Aspergillus niger</i>)	84 mg
Turmeric (<i>Curcuma longa</i>) (95% curcuminoids**)	7,5 mg
Grape seed (<i>Vitis vinifera</i>) (80% OPC)	20 mg
Green tea (<i>Camellia sinensis</i>) (75% EGCG; 3,5 mg/caps.)	14 mg
Lutein (from <i>Tagetes erecta</i>)	0,7 mg
Lycopene (from <i>Lycopersicon esculentum</i>)	1,7 mg

* NRV: Nutrient Reference Value in %
 ** provides curcumin I, demetoxicurcumin and bisdemetoxicurcumin

Recommended daily dose: 3 capsules one to two times daily with food. If you are taking medications, take this product a few hours before or after taking them.

Calcium triphosphate instead of bovine microcrystalline hydroxyapatite. Glucosamine of plant origin (from *Aspergillus niger*) instead of crustaceans.

References:

- Castelo-Branco, C., and J. Dávila Guardia. "Use of ossein-hydroxyapatite complex in the prevention of bone loss: a review." *Climacteric* 18.1 (2015): 29-37.
- Pines, A., et al. "Clinical trial of microcrystalline hydroxyapatite compound ('Ossopan') in the prevention of osteoporosis due to corticosteroid therapy." *Current medical research and opinion* 8.10 (1984): 734-742.
- Rüeggsegger, P., A. Keller, and M. A. Dambacher. "Comparison of the treatment effects of ossein-hydroxyapatite compound and calcium carbonate in osteoporotic females." *Osteoporosis international* 5.1 (1995): 30-34.
- Kamhi, Ellen. "Naturopathic Approaches to Preventing and Treating Osteoporosis." *Natural Medicine Journal* 2.11 (2010).
- Carpenter, Thomas O., et al. "A randomized controlled study of effects of dietary magnesium oxide supplementation on bone mineral content in healthy girls." *The Journal of Clinical Endocrinology & Metabolism* 91.12 (2006): 4866-4872.
- Rosenthal, Julie M., et al. "Dose-ranging study of lutein supplementation in persons aged 60 years or older." *Investigative ophthalmology & visual science* 47.12 (2006): 5227-5233.
- Devine, Amanda, et al. "Tea drinking is associated with benefits on bone density in older women." *The American journal of clinical nutrition* 86.4 (2007): 1243-1247.
- Henrotin, Yves, et al. "Decrease of a specific biomarker of collagen degradation in osteoarthritis, Coll2-1, by treatment with highly bioavailable curcumin during an exploratory clinical trial." *BMC complementary and alternative medicine* 14.1 (2014): 159.
- Lin, Yumei, et al. "Bone health nutraceuticals alter microarray mRNA gene expression: A randomized, parallel, open-label clinical study." *Phytotherapy* 23.1 (2016): 18-26.
- Corletto, F. "THERAPEUTICAL NOTES Female climacteric osteoporosis therapy with titrated horsetail (*Equisetum arvense*) extract plus calcium (osteosil calcium): randomized double blind study." *Minerva Ortopedica e Traumatologica* 50.5 (1999): 201-206.
- Prentice, Ross L., et al. "Health risks and benefits from calcium and vitamin D supplementation: Women's Health Initiative clinical trial and cohort study." *Osteoporosis International* 24.2 (2013): 567-580.
- Inaba, Naoko, Toshiro Sato, and Takatoshi Yamashita. "Low-Dose Daily Intake of Vitamin K2 (Menaquinone-7) Improves Osteocalcin -Carboxylation: A Double-Blind, Randomized Controlled Trials." *Journal of nutritional science and vitaminology* 61.6 (2015): 471-480.
- Whelan, Anne Marie, Tannis M. Jurgens, and Susan K. Bowles. "Natural health products in the prevention and treatment of osteoporosis: systematic review of randomized controlled trials." *Annals of Pharmacotherapy* 40.5 (2006): 836-849.

MENOPEACE

6 HIGHLY EFFECTIVE, SOY-FREE, STANDARDISED PLANT EXTRACTS FOR COMPREHENSIVE MENOPAUSAL SUPPORT



Product Datasheet

60 capsules Code FE0232

Nutritional information:	1 capsule
Dong quai ⁽¹⁾ (1% Z-ligustilide)	125 mg
Chaste tree ⁽¹⁾ (0,5% agnusides)	125 mg
Red clover ⁽¹⁾ (40% isoflavones; 25,2 mg AIE*)	75 mg
Dandelion ⁽¹⁾ (3% flavonoids)	63 mg
<i>Cimicifuga racemosa</i> ⁽¹⁾ (2,5 % triterpenes)	30 mg
<i>Dioscorea villosa</i> ⁽¹⁾ (16% diosgenin) (8:1)	25 mg

*AIE: Aglycone Isoflavone Equivalents

⁽¹⁾ standardised extracts

Recommended daily dose: 1–2 capsules daily. Use for a minimum of 3 months.

Specifications:

- » Formula rich in phytoestrogens that act as hormone regulators.
- » The components of this formula are suitable for helping relieve the various inconvenient discomforts associated with menopause (1-3).
- » The plants used in this formula are standardised extracts to ensure the effectiveness of the active ingredients.

DONG QUAI (*Angelica sinensis*)⁽⁴⁾

- » Female tonic
- » Dysmenorrhoea, general discomfort and bloating
- » Mood swings

CHASTE TREE (*Vitex agnus-castus*)⁽⁵⁻⁶⁾

- » Female hormonal balance
- » Breast pain (mastalgia), swelling, retention, anxiety, irritability, mood swings

BLACK COHOSH (*Cimicifuga racemosa*)⁽⁷⁻⁸⁾

- » Uterine cramps
- » Depression, anxiety, mood swings

RED CLOVER (*Trifolium pratense*)⁽⁹⁻¹⁰⁾

- » Oestrogen modulator
- » Greater contribution and more complete isoflavone profile compared to soy

DANDELION (*Taraxacum officinale*)⁽¹¹⁾

- » Fluid retention
- » Hepatic detoxification oestrogen elimination

DIOSCOREA VILLOSA OR MEXICAN WILD YAM⁽¹²⁾

- » Enhances the phytoestrogenic action of other plants



References:

1. Kronenberg, Fredi, and Adriane Fugh-Berman. "Complementary and alternative medicine for menopausal symptoms: a review of randomized, controlled trials." *Annals of internal medicine* 137.10 (2002): 805-813.
2. Mazarro Costa, Renata, et al. "Medicinal plants as alternative treatments for female sexual dysfunction: utopian vision or possible treatment in climacteric women?" *The journal of sexual medicine* 7.11 (2010): 3695-3714.
3. Rotem, Carmela, and Boris Kaplan. "Phyto-Female Complex for the relief of hot flashes, night sweats and quality of sleep: randomized, controlled, double-blind pilot study." *Gynecological endocrinology* 23.2 (2007): 117-122. Dong quai (*Angelica sinensis*)
4. Hirata, Janie D., et al. "Does dong quai have estrogenic effects in postmenopausal women? A double-blind, placebo-controlled trial." *Fertility and sterility* 68.6 (1997): 981-986. Sauzgatillo (*Vitex agnus-castus*):
5. Zamani, Mehrangiz, Nosrat Neghab, and Saadat Torabian. "Therapeutic effect of *Vitex agnus castus* in patients with premenstrual syndrome." *Acta Medica Iranica* 50.2 (2012): 101.
6. Momoeda, Mikio, et al. "Efficacy and safety of *Vitex agnus-castus* extract for treatment of premenstrual syndrome in Japanese patients: a prospective, open-label study." *Advances in therapy* 31.3 (2014): 362-373. *Cimicifuga racemosa*:
7. Frei-Kleiner, S., et al. "*Cimicifuga racemosa* dried ethanolic extract in menopausal disorders: a double-blind placebo-controlled clinical trial." *Maturitas* 51.4 (2005): 397-404.
8. Nappi, Rossella E., et al. "Efficacy of *Cimicifuga racemosa* on climacteric complaints: a randomized study versus low-dose transdermal estradiol." *Gynecological Endocrinology* 20.1 (2005): 30-35. Trébol rojo (*Trifolium pratense*):
9. Del Giorno, Cecilia, et al. "Effects of *Trifolium pratense* on the climacteric and sexual symptoms in postmenopausal women." *Rev Assoc Med Bras* 56.5 (2010): 558-562.
10. Booth, Nancy L., et al. "Clinical studies of red clover (*Trifolium pratense*) dietary supplements in menopause: a literature review." *Menopause* 13.2 (2006): 251-264. Diente de león (*Taraxacum officinale*)
11. Clare, Bevin A., Richard S. Conroy, and Kevin Spelman. "The diuretic effect in human subjects of an extract of *Taraxacum officinale* folium over a single day." *The Journal of Alternative and Complementary Medicine* 15.8 (2009): 929-934. *Dioscorea villosa*:
12. Komesaroff, P. A., et al. "Effects of wild yam extract on menopausal symptoms, lipids and sex hormones in healthy menopausal women." *Climacteric* 4.2 (2001): 144-150.

FOLATE ULTRA

CALCIUM L-METHYLFOLATE + VITAMIN B6 + VITAMIN B12

Folate intake is a critical factor in the prevention of neural tube defects, some forms of anaemia, cardiovascular disease and cognitive dysfunction.

Among the different chemical forms of folates, calcium L-methylfolate (5-methyltetrahydrofolate, 5-MTHF) is the most bio-available form.



60 tablets Code FE2037

[Product datasheet](#)

Nutritional information:	1 tablet
Folate (calcium-L-methylfolate)	1 000 µg (500%*)
Vitamin B ₆ (from 2 mg pyridoxal 5'-phosphate)	2 mg (143%*)
Vitamin B ₁₂ (methylcobalamin)	2,8 µg (112%*)

*NRV: Nutrient Reference Value in %

Recommended daily dose: 1 tablet daily.

- » Includes **Vitamins B6** and **B12** which act as cofactors in metabolic reactions in the body.

5-MTHF HAS SIGNIFICANT ADVANTAGES OVER FOLIC ACID:

- » **Better absorption** even when gastrointestinal pH is altered
- » **High bioavailability** unaffected by metabolic disturbances
- » **B12 deficiency** reduces the masking of B12 deficiency symptoms
- » **Low interaction** with dihydrofolate reductase inhibiting medications
- » **Overcomes metabolic defects** caused by MTHFR gene polymorphism
- » **Prevents the negative effects** of high concentrations of unconverted folic acid in peripheral circulation.

References:

1. Scaglione F. and Panzavolta G. Folate, folic acid and 5-methyltetrahydrofolate are not the same thing. Xenobiotica, 2014; 44(5): 480-488.

IRON BISGLYCINATE COMPLEX

HIGH BIOAVAILABILITY WITHOUT GASTROINTESTINAL DISCOMFORT



30 capsules FE2036

[WEB INFORMATION](#)

Nutritional information:	1 capsule
Iron (ferrous bisglycinate)	35 mg (250 %*)
Copper (cupric citrate)	1 mg (100 %*)
Thiamin (vit. B1)	4,5 mg (409 %*)
Riboflavin (vit. B2)	3,75 mg (268 %*)
Niacin (vit. B3)	5 mg (31 %*)
Vitamin B6	3,42 mg (244 %*)
Folate	1 000 µg (500 %*)
Vitamin B12	1 000 µg (40 000 %*)
Vitamin C	75 mg (94 %*)

*NRV: Nutrient Reference Value in %

Recommended daily dose: 1 capsule daily with food. If you are taking medication, take this product a few hours before or after them.

- » Ferrous bisglycinate has an absorption rate up to 3 times higher than other forms of iron, as it is absorbed through an alternative pathway within the intestinal mucosa. In this way, it does not interfere with other forms of iron in the diet.
- » It does not cause the common side effects of other iron supplements such as stomach pain, constipation, diarrhoea or cramps.
- » **With B-complex vitamins** essential for the formation of red blood cells
- » **With Vitamin C and copper** which improve the absorption of iron

- » A recent study found that those with low iron levels were more prone to strokes compared to those with normal iron levels.
- » This and other studies suggest an association between iron deficiency and the risk of ischaemic stroke in both adults and children⁽²⁻⁴⁾.

References:

1. Shovlin CL, Chamali B, Santhirapala V, Livesey JA, Angus G, Manning R, et al. Ischaemic Strokes in Patients with Pulmonary Arteriovenous Malformation and Hereditary Hemorrhagic Telangiectasia: Associations with Iron Deficiency and Platelets. PLoS One. 2014; 9(2): e88812.
2. Maquire JL, deVeber G, Parkin PC. Association between Iron Deficiency Anemia and Stroke in Young Children. Pediatrics. 2007; 120:1053.
3. Chang YL, Hung SH, Ling W, Lin HC, Li HC, Chung SD. Association between Ischemic Stroke and Iron-Deficiency Anemia: A population-Based Study. PLoS One. 2013; 8(12): e82952.
4. Dubyl MD, Card RT, Whiting SJ, oyle CA, Zlotkin SH, Paterson PG. Iron deficiency anemia prevalence at first stroke or transient ischemic attack. Can J Neurol Sci. 2012; 39(2): 189-95.

GRAPEFRUITSEED & CRANBERRY

500MG GRAPE SEED (95% PROANTHOCYANIDINS) + 100MG CRANBERRY (107:1) X CAPSULE



Product Datasheet

60 capsules Code FE1225

Nutritional information:	1 capsule
Grapeseed (95% proanthocyanidins)	500 mg
Cranberry (107:1)	100 mg

Recommended daily dose: 1-2 capsules daily.

Specifications:

- » Prevention of cardiovascular diseases.
- » Treatment and prevention of venous and capillary disorders.
- » Inflammation of joints and tissue of the stomach and intestines.
- » Strengthening of the immune system.
- » Retinal disorders.
- » Enhancing the anti-ageing action of other cosmetics.



ANTIOXIDANT

- » High antioxidant power due to its high OPC content, protecting the body from free radicals (2,3).
- » Prevents ageing and premature deterioration of cells, tissues and organs (2).

ANTI-INFLAMMATORY & ANTIBACTERIAL

- » It has anti-inflammatory action on stomach, intestinal and joint tissues (5).
- » Helps with arthritis, dermatitis, skin problems, sinusitis, colitis, urinary tract infection and asthma (5). It fights bacteria such as *E. coli* (6,12).

RETINOPATHY & ANTI-AGEING

- » Improves vision due to its high vitamin E content (13).
- » It helps to make skin healthier due to its resveratrol content and its ability to stimulate collagen production, making the skin more elastic (4).

VASOPROTECTIVE & CARDIOPROTECTIVE

- » It acts directly on the capillaries, keeping them strong and healthy (7-10).
- » It prevents hardening of the arteries and prevents thrombus formation (8,9).
- » It prevents arteriosclerosis and hypertension (10,11).



References:

1. Howell AB. Cranberry proanthocyanidins and the maintenance of urinary tract health. *Crit. Rev. Food Sci. Nutr.* 2002; 42:273-278
2. Koga T, et al. Increase of antioxidative potential of rat plasma by oral administration of proanthocyanidin-rich extract from grape seeds. *Journal of agricultural and food chemistry.* 1999;47(5):1892-1897
3. Bagchi D, et al. Oxygen free radical scavenging abilities of vitamins C and E, and a grape seed proanthocyanidin extract in vitro. *Research Communications in Molecular Pathology and Pharmacology.* 1997;95(2):179-189
4. Deckert V, et al. Prevention of LDL -to-copherol consumption, cholesterol oxidation, and vascular endothelium dysfunction by polyphenolic compounds from red wine. *Atherosclerosis.* 2002;165(1):41-50
5. Li W, et al. Anti-inflammatory effect and mechanism of proanthocyanidins from grape seeds. *Acta Pharmacologica Sinica.* 2000;22(12):1117-1120
6. Itzhak O, et al. Anti-Escherichia coli Adhesion Activity of Cranberry and Blueberry Juices. *N Engl J Med.* 1991;324:1599
7. Fitzpatrick DF, et al. Vasodilating pro-cyanidins derived from grape seeds. *Annals of the New York Academy of Sciences.* 2002; 957(1):78-89
8. Fitzpatrick DF, et al. Isolation and characterization of endothelium-dependent vasorelaxing compounds from grape seeds. *Journal of Agricultural and Food Chemistry.* 2000;48(12):6384-6390
9. Fitzpatrick DF, et al. Vasodilating pro-cyanidins derived from grape seeds. *Annals of the New York Academy of Sciences.* 2002;957(1):78-89
10. Aldini G, et al. Procyanidins from grape seeds protect endothelial cells from peroxynitrite damage and enhance endothelium-dependent relaxation in human artery: new evidences for cardio-protection. *Life sciences.* 2003;73(22):2883-2898
11. Da Luz PL, et al. The effect of red wine on experimental atherosclerosis: lipid-independent protection. *Experimental and molecular pathology.* 1999;65(3):150-159
12. Bodet C, et al. Anti-inflammatory Activity of a High-molecular-weight Cranberry Fraction on Macrophages Stimulated by Lipopolysaccharides from Periodontopathogens. *Journal of dental research.* 2006; 85(3):235-239
13. Zafra-Stone S, et al. Health Benefits of Edible Berry Anthocyanins: Novel Antioxidant and Anti-Angiogenic Properties. *Antioxidant Measurement and Applications.* 2007;22:337-351
14. Howell AB. Bioactive compounds in cranberries and their role in prevention of urinary tract infections. *Molecular nutrition & food research* 2007;51(6):732-737
15. Chi-Hua V, et al. Antibacterial effects of American cranberry (*Vaccinium macrocarpon*) concentrate on foodborne pathogens. *LWT-Food Science and Technology.* 2008;41(10):1834-1841
16. Katalinic V, et al. Polyphenolic profile, antioxidant properties and antimicrobial activity of grape skin extracts of 14 *Vitis vinifera* varieties grown in Dalmatia (Croatia). *Food Chemistry.* 2010;119(2):715-723.

Notes





What Makes New Roots Herbal Unique?

- » At New Roots Herbal our complexes are formulated by naturopathic doctors, scientists and herbalists to address imbalances in a wide-range of chronic and acute health conditions.
- » We are committed to providing pure and high-quality nutrients and powerful botanical herbs that work synergistically to support the body.
- » The products are tested for potency and purity (including all contaminants, such as heavy metals and PCBs) through the production process to the final product. This we guarantee you!
- » Our aim is to support and educate practitioners on the powerful synergetic properties of herbs and nutraceutical blends. Our Nutritional Therapist and online resources offer on-hand advice and guidance so you can provide the best programmes and service to your clients.

